



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

head, front of thorax, base of abdomen, sides and tip of abdomen white, the rest brown and blackish; abdominal tufts dark brown. Expanse, 64 mm.

One female, Mexico City, Mexico, June, 1907 (R. Müller, No. 1034).

Type. — No. 10456, U. S. National Museum.

The specimen agrees almost exactly with Felder and Rogenhofer's figure of *Amphidasys cladonia* (Plate CXXV, Fig. 13), the marginal spottings being only somewhat more extensive. Were it not that the locality for *cladonia* is given as "Silhet" I should think that the species was before me.

EUCYMATOGE RECTILINEATA, A NEW GEOMETRID MOTH FROM COLORADO.

By GEO. W. TAYLOR,

WELLINGTON, B. C., CANADA.

This species belongs to the group containing in our fauna *E. intestinata*, *gillettei* and *vitalbata*, and to which the generic name *Phibalapteryx* Stephens, used to be applied. It resembles *vitalbata* more nearly than it does *intestinata*, but it is smaller, and the colors are less sharply contrasting. It may be separated at once from any of the three species mentioned by the *straightness* of the extra discal line.

I have had a single specimen of this species in my cabinet without name for a considerable time. The discovery of a second specimen, exactly similar, in the collection of the U. S. National Museum, determines me to describe the form as new.

***Eucymatoge rectilineata*, new species.**

Expanse 25 mm. Prevailing color of wings, warm chocolate brown with the costal and basal areas and the hinder portion of the submarginal band lighter as in *E. vitalbata*. The hind wings are uniformly of this lighter color and not blackish in the median band as in the last named species. The fore wings are crossed by numerous lines as in the other species in the group, but in *rectilineata* the extra discal line runs in an almost straight line from vein 3 to the center of the inner margin and not in a series of scallops. The hind wings are clearer than in the other species, and the lines are pale on a darker ground.

Described from two specimens, one bearing label "Colorado, Las Pinas, 30 May," is in my own collection and the second labelled "Durango, Col., July 8-15," is in the U. S. National Museum and bears their type No. 10371.